

On the measurement probability of quantum phases

Thomas Schürmann

Jülich Supercomputing Centre, Jülich Research Centre, 52425 Jülich, Germany

We consider the probability by which quantum phase measurements of a given precision can be done successfully. The *least upper bound* of this probability is derived and the associated optimal state vectors are determined. The probability bound represents an unique and continuous transition between macroscopic and microscopic measurement precisions.

PACS numbers: 42.50.-p, 03.65.Ta

The classical picture for the evolution of a single-mode electromagnetic field is simply determined by an amplitude (specifying the strength of the field) and a phase (specifying the zeros of the field). On the other hand, the concept of electromagnetic phase as an observable quantity is a long-standing problem of quantum optics and it has been the question whether there exists a phase observable that is canonically conjugate to the number observable for a single-mode field. The quantum mechanical description of phase was first considered by London [1] and Dirac [2]. An obvious way of defining an operator for the phase is by polar decomposition of the photon annihilation operator $\hat{a} = e^{i\hat{\phi}}\sqrt{\hat{N}}$. The phase operator $\hat{\phi}$ defined in this way is equivalent to that considered by Dirac [2], who obtained the commutator $[\hat{\phi}, \hat{N}] = i$ by employing the correspondence between commutators and classical Poisson bracket. Formally, this would imply the uncertainty relation

$$\sigma_{\phi}\sigma_N \geq \frac{1}{2} \quad (1)$$

with σ_{ϕ} and σ_N are the standard deviations of ϕ and N . The difficulties of Dirac's approach were clearly pointed out by Susskind and Glogower [3]. Firstly, the relation (1) would imply that a well-defined number state would have a phase standard deviation greater than 2π . This is a consequence of the fact that Dirac's commutator does not take account of the periodic nature of the phase. Furthermore, the exponential operator $e^{i\hat{\phi}}$ derived from this approach is not unitary and thus does not define a Hermitian operator. This is why it is often accepted that a well-behaved Hermitian phase operator does not exist [3, 4]. Therefore, arguments based on the Heisenberg relation (1) cannot hold in general.

Actually, the standard deviation offers a reasonable measure of the spread of values when the distribution in question is of a simple "single hump" type. In particular it is a very good characteristic for a Gaussian distribution since it measures directly the half-width of this distribution. However, when the distribution is not of a simple type (for example, has more than one hump) the standard deviation loses much of its usefulness as a measure of uncertainty.

The aim of the present contribution is to introduce the probability by which a successful phase measurements of

a given precision can be done. The *least upper bound* of this probability is determined and the corresponding (optimal) state vectors are computed.

In order to specify phase measurements, the probability distribution for the measurement result can be determined using *positive operator-valued measures*. This approach was first considered by Helstrom [5], and is also considered in [6, 7]. Precisely, let $\mathcal{H}$ be a complex separable Hilbert space, $(|n\rangle)_{n \geq 0}$ an orthonormal basis, and $N = \sum_{n=0}^{\infty} n|n\rangle\langle n|$ the associated number observable. If the phase density F_{ϕ} treats all phases equally, it should be invariant under phase translation, $e^{i\theta N}F_{\phi}e^{-i\theta N} = F_{\phi+\theta}$, generated by the number observable. In this case, the general form of F_{ϕ} is

$$F_{\phi} = \frac{1}{2\pi} \sum_{n,m=0}^{\infty} c_{nm} e^{i(n-m)\phi} |n\rangle\langle m| \quad (2)$$

where (c_{nm}) is the associated phase matrix. For the integral of the probability to equal 1, we must have

$$\int_{-\pi}^{\pi} F_{\phi} d\phi = 1. \quad (3)$$

Applying this to (2) above we find that [8]

$$\sum_{n=0}^{\infty} c_{nn} |n\rangle\langle n| = 1. \quad (4)$$

This means that the diagonal elements c_{nn} must all be equal to 1. The additional condition of positive definite probabilities, together with the above result means that all of the c_{nm} must have absolute values between 0 and 1. In general, real measurements will give smaller values of c_{nm} , and the closer these are to 1 the better the phase measurement is. In [9] it is shown that the additional condition that a number shifter does not alter the phase distribution gives $c_{nm} = 1$, corresponding to the canonical measure [9]

$$E_{\phi} = \frac{1}{2\pi} \sum_{n,m=0}^{\infty} e^{i(n-m)\phi} |n\rangle\langle m|. \quad (5)$$

An alternative derivation of this result is by using the maximum likelihood approach [6]. Note that (5) may be

expressed by $dE_\phi = |\phi\rangle\langle\phi| d\phi$, where

$$|\phi\rangle = \frac{1}{\sqrt{2\pi}} \sum_{n=0}^{\infty} e^{in\phi} |n\rangle \quad (6)$$

With reference to (5), we now define the precision $\Delta\alpha \in [0, 2\pi)$ of a phase measurement corresponding to the vicinity $A_\alpha = [\alpha - \frac{\Delta\alpha}{2}, \alpha + \frac{\Delta\alpha}{2})$ of any value $\alpha \in [-\pi, \pi)$. The probability of a phase measurement with $\phi \in A_\alpha$, made on a state described by a density operator $\hat{\rho}$, is given by

$$P_\alpha(\Delta\alpha) = \text{Tr}[\hat{\rho} E_\phi(A_\alpha)]. \quad (7)$$

On the other hand, the probability to measure a photon number $n \in B_k$ is given by

$$P_k(\Delta k) = \text{Tr}[\hat{\rho} E_N(B_k)] \quad (8)$$

where

$$E_N(B_k) = \sum_{n \in B_k} |n\rangle\langle n| \quad (9)$$

is the value of the spectral measure E_N on a set B_k of positive integers. In order to introduce the precision $\Delta k \in \mathbb{N}$ by which the photon number is measured, we note that the photon number is bounded from below. Therefore, we define the right-sided vicinity $B_k \subset \mathbb{N}$ of k by $B_k = \{k, k+1, \dots, k+\Delta k\}$. In this definition, the integer k is the smallest element of B_k . Alternative definitions are also possible but typically lead to necessary readjustments in certain cases for $k < \Delta k$. However, it will be seen later that our results are not dependent on the particular subscription. For technical purposes we apply the definition of the minimum integer-subscription. In the case of pure states $\hat{\rho} = |\psi\rangle\langle\psi|$, we obtain the probability

$$P_k(\Delta k) = \sum_{n=k}^{k+\Delta k} |\psi_n|^2 \quad (10)$$

and $\psi_n = \langle n|\psi\rangle$ is the number-space amplitude of ψ .

Now, we consider the case with an initial photon number preparation of a state ψ . A single mode is supposed to emerge in a state according to

$$\psi \rightarrow \psi' = \frac{E_N(B_k)\psi}{\|E_N(B_k)\psi\|}. \quad (11)$$

Afterwards, the number of photons is given with precision Δk . In this situation the uncertainty principle suggests that the more accurately the number is measured the greater is the perturbation of the phase of the outgoing state. The conditional probability $\mathcal{P}_{\alpha k}(\Delta\alpha | \Delta k; \psi)$ to measure phase $\phi \in A_\alpha$, on the state transformed by the initial number measurement, is given by

$$\mathcal{P}_{\alpha k}(\Delta\alpha | \Delta k; \psi) = \frac{\|E_\phi(A_\alpha)E_N(B_k)\psi\|^2}{\|E_N(B_k)\psi\|^2} \quad (12)$$

We now ask for the *least upper bound* λ_0 of the measurement probability (12) and we end up in a variation problem in Hilbert space with three degrees of freedom. For fixed precisions Δk and $\Delta\alpha$ we are searching for the supremum of (12) by variation of the parameters α , k and the state vector ψ of the photon. Actually this variation problem is translation and rotation invariant in Hilbert space and we can simply chose $k = 0$ and $\alpha = 0$ without loss of generality. After all, we have to consider the following expression

$$\lambda_0 = \sup_{\psi \in \mathcal{H} \setminus \{0\}} \mathcal{P}_{00}(\Delta\alpha | \Delta k; \psi) \quad (13)$$

and by using (5) and (9) we explicitly obtain to following expression

$$\mathcal{P}_{00}(\Delta\alpha | \Delta k; \psi) = \frac{\int_{-\Delta\alpha/2}^{\Delta\alpha/2} \left| \frac{1}{\sqrt{2\pi}} \sum_{n=0}^{\Delta k} \psi_n e^{in\phi} \right|^2 d\phi}{\sum_{n'=0}^{\Delta k} |\psi_{n'}|^2}. \quad (14)$$

Applying the Cauchy-Bunyakovsky inequality we obtain the following general upper bound

$$\mathcal{P}_{\alpha,k}(\Delta\alpha | \Delta k; \psi) \leq \frac{\Delta\alpha(\Delta k + 1)}{2\pi} \quad (15)$$

for every $\Delta\alpha \in [-\pi, \pi)$ and integer $\Delta k \geq 0$. In fig. 1, the set of impossible measurement processes is expressed by the grey shaded triangle. In order to reach even tighter




Figure 1: Maximum eigenvalue λ_0 for $\Delta k = 0, 1, 2, 3, \infty$ (from left to right). Phase measuring processes with probabilities larger than λ_0 do not exist.

bounds we explicitly computed the integral in (14) and applied certain trigonometric identities to obtain the following expression

$$\mathcal{P}_{00}(\Delta\alpha | \Delta k; \psi) = \sum_{n,m=0}^{\infty} G_{nm} \psi_n \psi_m^* \quad (16)$$

with normalization condition $\sum_{n=0}^{\Delta k} |\psi_n|^2 = 1$ and kernel [8]

$$G_{nm} = \frac{1}{\pi} \frac{\sin \frac{\Delta \alpha}{2} (n-m)}{n-m} \quad (17)$$

for $0 \leq n, m \leq \Delta k$, 0 otherwise. Obviously, G is self-adjoint and positive definite. According to (13) and (16), the least upper bound is given by the operator norm of G , i.e.

$$\lambda_0 = \|G\|, \quad (18)$$

and this norm is identical to the largest eigenvalue of G . In order to obtain the eigenvalues of G we have to solve the following linear equation for $n = 0, 1, \dots, \Delta k$

$$\sum_{m=0}^{\Delta k} G_{nm} \psi_m^{(s)} = \lambda_s \psi_n^{(s)} \quad (19)$$

for $s = 0, 1, \dots, \Delta k$. This type of eigenvalue problem has been extensively discussed in [11] (see also references therein). All eigenvalues are distinct, positive and may be ordered so that $1 > \lambda_0 > \lambda_1 > \dots > \lambda_{\Delta k}$. In the non-trivial case of $\Delta k > 0$ we computed λ_0 numerically. For increasing values of Δk , the corresponding bounds approach very fast to the asymptotic case $\Delta k \rightarrow \infty$, see fig. 1 (right most continuous line). For the computation of the asymptotic case we introduced the equidistant decomposition $q_m = \frac{m}{\Delta k+1}$, $m = 0, 1, \dots, \Delta k$ with increment $\delta q_m = q_{m+1} - q_m$. After substitution into (19) and a few algebraic manipulations, the discrete eigenvalue problem approaches to the following homogeneous Fredholm integral equation of the first kind

$$\frac{1}{\pi} \int_{-1}^1 \frac{\sin(\frac{\pi}{2} \xi (z-z'))}{z-z'} \varphi^{(\nu)}(z') dz' = \tilde{\lambda}_\nu(\xi) \varphi^{(\nu)}(z) \quad (20)$$

with $|z| \leq 1$ and

$$\xi = \frac{\Delta \alpha (\Delta k + 1)}{2\pi}. \quad (21)$$

From standard theory we know that there are solutions in $L^2([-1, 1])$ only for a discrete set of eigenvalues, say $\tilde{\lambda}_0 \geq \tilde{\lambda}_1 \geq \dots$ and that as $\nu \rightarrow \infty$, $\tilde{\lambda}_\nu \rightarrow 0$. It should be noted that the eigenvalues explicitly depend on the parameter ξ and corresponding to each eigenvalue there is a unique (up to normalization) solution $\varphi^{(\nu)}(z) = S_{0\nu}(\pi\xi/2, z)$ called *angular prolate spheroidal wave function* [10, 11]. They are continuous functions of ξ for $\xi \geq 0$, and are orthogonal in $(-1, 1)$. Moreover, they are complete in $L^2([-1, 1])$. The corresponding eigenvalues are related to a second set of functions called *radial prolate spheroidal functions*, which differ from the angular functions only by a real scale factor [11]. Applying the notation of [10, 11], these eigenvalues are

$$\tilde{\lambda}_\nu(\xi) = \xi \left[R_{0\nu}^{(1)}(\pi\xi/2, 1) \right]^2 \quad (22)$$

with $\nu = 0, 1, 2, \dots$. The properties of the discrete eigenvalue spectrum is discussed in [12]. Here, we are mainly interested in the properties of the largest eigenvalue $\tilde{\lambda}_0(\xi)$. It is monotonically increasing and approaches 1 exponentially in ξ . For small values of ξ there is the asymptotic behavior $\tilde{\lambda}_0(\xi) \sim \xi$.

-
- [1] F. London, Z. Phys. **37** (1926) 915; **40** (1927) 193.
 - [2] P. Dirac, Proc. Roy. Soc. London A **114** (1927) 243.
 - [3] L. Susskind and J. Glogower, Physics **1** (1964) 49.
 - [4] P. Carruthers and M. M. Nieto, Rev. Mod. Phys. **40** (1968) 411.
 - [5] C. W. Helstrom, *Quantum Detection and Estimation Theory* (Academic Press, New York, 1976).
 - [6] J. H. Shapiro, S. R. Shepard and N. C. Wong, Phys. Rev. Lett. **62** (1989) 2377.
 - [7] J. H. Shapiro and S. R. Shepard, Phys. Rev. A **43** (1991) 3795.
 - [8] It is understood here, that when $n = m$ the expression $\frac{\sin(\frac{\Delta \alpha}{2} (n-m))}{n-m}$ has the value $\frac{\Delta \alpha}{2}$.
 - [9] U. Leonhardt, J. A. Vaccaro, B. Böhmer and H. Paul, Phys. Rev. A **51** (1995) 84.
 - [10] M. Abramowitz and I. Stegun, *Handbook of Mathematical Functions*, (Dover, New York, 1965).
 - [11] D. Slepian, Bell Syst. Tech. Jour. **57** (1978) 1371.
 - [12] H. J. Landau, Trans. Amer. Math. Soc. **115** (1965) 242.